

On the Homothety Conjecture ^{*}

Elisabeth M. Werner [†]

Deping Ye [‡]

Abstract

Let K be a convex body in $\mathbb{R}^n$ and $\delta > 0$. The homothety conjecture asks: Does $K_\delta = cK$ imply that K is an ellipsoid? Here K_δ is the (convex) floating body and c is a constant depending on δ only. In this paper we prove that the homothety conjecture holds true in the class of the convex bodies B_p^n , $1 \leq p \leq \infty$, the unit balls of l_p^n ; namely, we show that $(B_p^n)_\delta = cB_p^n$ if and only if $p = 2$. We also show that the homothety conjecture is true for a general convex body K if δ is small enough. This improves earlier results by Schütt and Werner [16] and Stancu [20].

1 Introduction

Floating bodies appear in many contexts and have been widely studied (see e.g. [1, 2, 4, 5, 6, 8, 13, 14, 15, 19, 20, 21, 23, 26]). The homothety conjecture is among the problems related to floating bodies that was open for a long time. It asks:

Does K have to be an ellipsoid, if K is homothetic to K_δ for some $\delta > 0$?

In [16], Schütt and Werner obtained a (partial) positive solution to this conjecture. They showed that if there is a sequence $\delta_k \rightarrow 0$ such that K_{δ_k} is homothetic to K for all $k \in \mathbb{N}$ (with respect to the same center of homothety), then K is an ellipsoid.

Stancu [20] (see also [19]) proved that if for a convex body K with boundary of class C_+^2 there exists a positive number $\delta(K)$ such that K_δ is homothetic to K

^{*}Keywords: convex floating body, homothety problem. 2000 Mathematics Subject Classification: 52A20, 53A15

[†]Partially supported by an NSF grant, a FRG-NSF grant and a BSF grant.

[‡]Support by NSF-FRG DMS: 0652571.

for some $\delta < \delta(K)$, then and only then K is an ellipsoid.

These results are not completely satisfactory for different reasons. The first one requires a sequence K_{δ_k} of convex bodies to be homothetic to the body K . The second one needs smoothness assumptions on ∂K and, in addition, δ has to be sufficiently small, but no estimates are given how small. Moreover, these results do not work even in the case of very basic convex bodies, such as B_p^n , $1 \leq p \leq \infty$, the unit balls of l_p^n .

It should also be noted in this context, that Milman and Pajor [9] showed that if K is a symmetric convex body, then for large δ , the floating body K_δ is always uniformly, up to a factor $C(\delta)$, isomorphic to an ellipsoid, namely to the dual of the Binet ellipsoid from classical mechanics.

In this paper, we give a positive solution of the homothety conjecture. Namely we prove

Theorem 2 *Let K be a convex body in $\mathbb{R}^n$. There exists a positive number $\delta(K)$, such that K_δ is homothetic to K for some $\delta \leq \delta(K)$, if and only if K is an ellipsoid.*

Our proof is different from Stancu's proof. No smoothness assumptions are required. In fact, the main ingredient in our proof is to show that the homothety assumption implies that ∂K is C_+^2 . This is done in Lemmas 2 and 4.

Our proof of Theorem 2 gives a possibility to estimate the threshold $\delta(K)$ for convex bodies K in $\mathbb{R}^n$ that have sufficiently smooth boundary. This is done in Section 5.

We also show the following theorem, which gives a positive solution of the homothety conjecture for the B_p^n balls, $1 \leq p \leq \infty$, and their affine images without any requirements on the size of δ .

Theorem 1 *Let B_p^n , $1 \leq p \leq \infty$ be the unit ball of l_p^n . Let $0 < \delta < \frac{|B_p^n|}{2}$. Then $(B_p^n)_\delta = cB_p^n$ for some $0 < c < 1$ if and only if $p = 2$.*

Throughout the paper we will use the following notations. For $1 \leq p \leq \infty$,

$B_p^n = \{x \in \mathbb{R}^n : \|x\|_p \leq 1\}$ is the unit ball of $l_p^n = (\mathbb{R}^n, \|\cdot\|_p)$ where for $1 \leq p < \infty$

$$\|x\|_p = \left(\sum_{i=1}^n |x_i|^p \right)^{\frac{1}{p}},$$

and for $p = \infty$

$$\|x\|_\infty = \max_{1 \leq i \leq n} |x_i|.$$

Let $u \in S^{n-1} = \partial B_2^n$, the boundary of the Euclidean unit ball B_2^n . Then $H(x, u) = \{y \in \mathbb{R}^n, \langle y, u \rangle = \langle x, u \rangle\}$ is the hyperplane through x with outer unit normal vector u . The two half-spaces generated by $H(x, u)$ are $H^-(x, u) = \{y \in \mathbb{R}^n, \langle y, u \rangle \geq \langle x, u \rangle\}$ and $H^+(x, u) = \{y \in \mathbb{R}^n, \langle y, u \rangle \leq \langle x, u \rangle\}$. $\|\cdot\|$ is the Euclidean norm on $\mathbb{R}^n$.

For a convex body K in $\mathbb{R}^n$ and $x \in \partial K$, the boundary of K , $N_K(x)$ denotes the outer unit normal vector to K and $\kappa_K(x)$ the Gauss curvature of K at x . $N_K(x)$ exists almost everywhere (see[11]). $\text{int}(K)$ is the interior of K . We write $|K|$ or $\text{vol}_n(K)$ for the volume of K . We say that K is in C_+^2 , if ∂K is C^2 and has everywhere strictly positive Gauss curvature. Without loss of generality, we will assume throughout the paper that, unless specified otherwise, 0 is the center of gravity of K and that K homothetic to K_δ is meant with 0 as the center of homothety.

The paper is organized as follows. In Section 2 we provide background and prove some properties of the convex floating body that are needed in the next sections. In Section 3 we prove that the homothety conjecture holds true for B_p^n , $1 \leq p \leq \infty$. In Section 4 we prove Theorem 2. Moreover, we give (partial) positive solutions of a generalized homothety conjecture. In Section 5 we give estimates on the threshold $\delta(K)$.

2 The convex floating body and homothety

Let K be a convex body with $0 \in \text{int}(K)$ and δ be a positive number such that $\delta < \frac{|K|}{2}$. The convex floating body is defined as follows [15].

Definition 1 [15] *Let K be a convex body in $\mathbb{R}^n$. The convex floating body K_δ is the intersection of all halfspaces H^+ whose defining hyperplanes H cut off a set of*

volume at most δ from K ,

$$K_\delta = \bigcap_{|H^- \cap K| \leq \delta} H^+.$$

Clearly, $K_0 = K$ and $K_\delta \subset K$ for all $\delta \geq 0$. Moreover, the convex floating body has the following property: for all (invertible) affine maps T on $\mathbb{R}^n$ and for all $\delta > 0$

$$(TK)_\delta = T \left(K_{\frac{\delta}{|\det(T)|}} \right). \quad (1)$$

Here $|\det(T)|$ is the absolute value of the determinant of T . In particular, for an affine map T with $|\det(T)| = 1$, $(TK)_\delta = T(K_\delta)$ for all $\delta \geq 0$.

An ellipsoid $\mathcal{E}$ is the affine image of B_2^n , $\mathcal{E} = T(B_2^n)$, for some invertible affine map T on $\mathbb{R}^n$. It is easy to see that $(B_2^n)_\delta = cB_2^n$ for all $\delta \geq 0$ and for a constant $c = c(\delta) < 1$ depending on δ only. Hence one gets with (1) that for all ellipsoids $\mathcal{E}$

$$\mathcal{E}_\delta = (T(B_2^n))_\delta = T \left((B_2^n)_{\frac{\delta}{|\det(T)|}} \right) = T \left(c \left(\frac{\delta}{|\det(T)|} \right) B_2^n \right) = c \left(\frac{\delta}{|\det(T)|} \right) \mathcal{E}$$

for all $\delta \geq 0$ and for some constant $c \left(\frac{\delta}{|\det(T)|} \right) < 1$. In other words, if the homothety conjecture holds true, then $K_\delta = cK$ for some $\delta > 0$ and some constant $0 < c < 1$ if and only if K is an ellipsoid in $\mathbb{R}^n$.

Now we make some general observations concerning homothety of K with one of its convex floating bodies. First, only a strictly convex body can be homothetic to one of its convex floating bodies. This is a consequence of the flowing lemma proved in [16].

Lemma 1 [16] *Let K be a convex body in $\mathbb{R}^n$, and let $0 < \delta < |K|/2$. Then K_δ is strictly convex.*

Thus, in particular, no polytope can be homothetic to one of its floating bodies. The next lemma and its proof is almost identical to Lemma 3 of [16]. We give the proof for completeness.

Lemma 2 *Let K be a convex body in $\mathbb{R}^n$, and let $\delta > 0$. If K_δ is homothetic to K , then ∂K is of class C^1 .*

Proof. Suppose that ∂K is not of class C^1 . Then there is $x_0 \in \partial K$ so that ∂K has two different supporting hyperplanes, H_1 and H_2 , passing through x_0 . We may assume that there are two sequences $(x_k^1)_{k \in \mathbb{N}}$ and $(x_k^2)_{k \in \mathbb{N}}$ on ∂K converging to x_0 so that we have:

- (i) the supporting hyperplanes H_k^i through x_k^i , $k \in \mathbb{N}$, $i = 1, 2$ are unique;
- (ii) the hyperplanes H_k^i converge to H_i , $i = 1, 2$;
- (iii) $\lim_{k \rightarrow \infty} N_K(x_k^i)$ is orthogonal to H_i , $i = 1, 2$.

See the proof of Lemma 3 in [16] for the construction of these sequences. We choose a coordinate system such that $x_0 = 0$,

$$H_1 = \{x \in \mathbb{R}^n : x(n) = ax(n-1)\},$$

and

$$H_2 = \{x \in \mathbb{R}^n : x(n) = bx(n-1)\},$$

with $b < a$ where $x(l)$ denotes the l -th coordinate of the vector $x \in \mathbb{R}^n$. Let x_δ be the point on ∂K_δ that corresponds to x_0 by homothety. We can assume that in the chosen coordinate system $x_\delta = (0, \dots, 0, x_\delta(n))$ with $x_\delta(n) > 0$. Let

$$H = \left\{ x + x_\delta \in \mathbb{R}^n : x(n) = \frac{a+b}{2}x(n-1) \right\}.$$

By homothety, one sees that H is a support hyperplane of ∂K_δ . For $\alpha, \beta \in \mathbb{R}$, let

$$M(\alpha, \beta) = \{x \in \mathbb{R}^n : x \in K \cap H^-, x(n-1) = \alpha, x(n) = \beta\},$$

and let M^* be such a set for which the $(n-2)$ -dimensional volume is maximal

$$\text{vol}_{n-2}(M^*) = \max_{(\alpha, \beta) \in \mathbb{R}^2} \text{vol}_{n-2}(M(\alpha, \beta)).$$

We consider the set in the $(x(n-1), x(n))$ -plane that consists of all points (α, β) so that $M(\alpha, \beta) \neq \emptyset$. This set is contained in the triangle T bounded by the lines $x(n) = ax(n-1)$, $x(n) = bx(n-1)$, and $x(n) = \frac{a+b}{2}x(n-1) + x_\delta(n)$. Therefore

$$\begin{aligned} \delta \leq |K \cap H^-| &= \int_{\mathbb{R}^2} M(\alpha, \beta) d(\alpha, \beta) \leq \text{vol}_{n-2}(M^*) \text{vol}_2(T) \\ &= \text{vol}_{n-2}(M^*) \frac{2|x_\delta(n)|^2}{a-b}. \end{aligned} \tag{2}$$

Now we consider the sets $K \cap (x_\delta + H_i)^-$, $i = 1, 2$. It follows from (i), (ii) and Lemma 2 of [16] that

$$|K \cap (x_\delta + H_i)^-| = \delta, \quad i = 1, 2.$$

This is because $x_\delta + H_i, i = 1, 2$ are tangent hyperplanes of K_δ . Let $\varepsilon > 0$ be given. By (iii) we may choose x_1 and x_2 in ∂K such that for $i = 1, 2$

$$\begin{aligned} \left(|x_i(n-1) - P_{H_i}(x_i)(n-1)|^2 + |x_i(n) - P_{H_i}(x_i)(n)|^2 \right)^{\frac{1}{2}} \leq \\ \varepsilon \left(|P_{H_i}(x_i)(n-1)|^2 + |P_{H_i}(x_i)(n)|^2 \right)^{\frac{1}{2}}, \end{aligned} \quad (3)$$

where P_{H_i} denotes the orthogonal projection onto $H_i, i = 1, 2$. As $K_\delta \subset K$, we can choose x_1 and x_2 in ∂K such that for $i = 1, 2$, $P_{H_i+x_\delta}(x_i) \in K$ and (3) still holds.

Moreover, as K is strictly convex, by Lemma 1 and homothety, we have that for $i = 1, 2$

$$\left(|x_i(n-1) - P_{H_i}(x_i)(n-1)|^2 + |x_i(n) - P_{H_i}(x_i)(n)|^2 \right)^{\frac{1}{2}} > 0.$$

It follows that there is a constant $c > 0$ that depends only on K so that

$$x_\delta(n) \leq c \left(|x_i(n-1) - P_{H_i}(x_i)(n-1)|^2 + |x_i(n) - P_{H_i}(x_i)(n)|^2 \right)^{\frac{1}{2}}. \quad (4)$$

For $i = 1$ or $i = 2$ the convex hull

$$[M^*, [x_i, P_{H_i+x_\delta}(x_i)]] \subset K \cap (x_\delta + H_i)^-.$$

We may assume that $i = 1$. Then

$$\begin{aligned} \delta = |K \cap (x_\delta + H_1)^-| &\geq \left| [M^*, [x_1, P_{H_1+x_\delta}(x_1)]] \right| \\ &\geq \frac{1}{n^2} \text{vol}_{n-2}(M^*) d(x_1, H_{M^*}) d(P_{H_1+x_\delta}(x_1), \tilde{H}), \end{aligned} \quad (5)$$

where H_{M^*} is the $(n-2)$ -dimensional flat containing M^* and $\tilde{H}$ is the plane containing M^* and x_1 . If ε is sufficiently small, we obtain by elementary computation

$$d(x_1, H_{M^*}) \geq c_1 \left(|P_{H_1}(x_1)(n-1)|^2 + |P_{H_1}(x_1)(n)|^2 \right)^{\frac{1}{2}},$$

and

$$d(P_{H_1+x_\delta}(x_1), \tilde{H}) \geq c_2 |x_\delta(n)|,$$

where c_1, c_2 depend only on K . Therefore we obtain by (2) and (5)

$$\frac{2|x_\delta(n)|}{a-b} \geq \frac{c_1 c_2}{n^2} \left(|P_{H_1}(x_1)(n-1)|^2 + |P_{H_1}(x_1)(n)|^2 \right)^{\frac{1}{2}}.$$

By (3) we get with a new constant $c > 0$

$$|x_\delta(n)| \geq \frac{c}{\varepsilon} \left(|x_1(n-1) - P_{H_1}(x_1)(n-1)|^2 + |x_1(n) - P_{H_1}(x_1)(n)|^2 \right)^{\frac{1}{2}}.$$

If we choose ε sufficiently small we get a contradiction to (4).

Remark 1 *If K is strictly convex and C^1 , K_δ need not be C^1 .*

To see that, consider a “rounded simplex” S in $\mathbb{R}^2$ whose vertices have been “rounded” by putting a small εB_2^n ball at each vertex and whose edges have been made strictly convex by replacing them by arcs of Euclidean balls RB_2^n with R very large. Then there are points of non-differentiability on that parts of ∂S_δ that face the “edges” of S . Hence homothety is crucial in the previous lemma.

Note however that if K is strictly convex, C^1 and, in addition, symmetric, then K_δ is C_+^2 . This was shown in [6].

The following symmetric matrix is closely related to the curvature of the floating body [4] (see also [16]). For $x \in \text{int}(K)$, $\xi \in S^{n-1}$, and an orthonormal coordinate system in the plane $H(x, \xi)$ with origin x , let $l(\eta)$ be the line through x with direction η and $y = l(\eta) \cap \partial K$. Let $r(\eta) = \|x - y\|$ and $\beta(\eta)$ be the angle between $l(\eta)$ and a tangent line to ∂K through y whose orthogonal projection onto $H(x, \xi)$ is $l(\eta)$. Define for $1 \leq i, j \leq n-1$

$$Q(i, j) = \frac{1}{|K \cap H(x, \xi)|} \int_{S^{n-2}} \eta_i \eta_j r^n(\eta) \cot(\beta(\eta)) d\sigma_{n-2}(\eta), \quad (6)$$

where σ_{n-2} is the surface measure on S^{n-2} . For $x_\delta \in \partial K_\delta$, we will use $H(x_\delta, \delta)$ to denote the hyperplane through x_δ cutting off a set of volume δ from K . Such a hyperplane always exists by Lemma 2 of [16].

Then we have the following lemma [4] (see also [16])

Lemma 3 [4] *Let K be a convex body of class C^1 and let $\delta > 0$. Suppose that for every $x_\delta \in \partial K_\delta$ and every $H(x_\delta, \delta)$ the matrix Q is positive definite. Then ∂K_δ is of class C^2 . Moreover, the Gauss curvature of K_δ at $x_\delta \in \partial K_\delta$ can be calculated by*

$$\frac{1}{\kappa_{K_\delta}(x_\delta)} = \det(Q),$$

and hence, K_δ is of class C_+^2 .

Also the next lemma and its proof is almost identical to Lemma 6 of [16]. Again, we give the proof for completeness. We will also use the following:

For every $x \in \partial K$, we define $\rho(x)$ to be the radius of the largest Euclidean ball that is contained in K and whose center lies on the line through x with direction $N_K(x)$,

$$\{x + t N_K(x) : t \in \mathbb{R}\}.$$

$\rho(x)$ is well defined because ∂K is of class C^1 . Let $t(x) \geq \rho(x)$ be such that $x - t(x) N_K(x)$ is a center of a ball with radius $\rho(x)$ that is contained in K . $\rho(x)$ is a continuous, strictly positive function on ∂K because ∂K is of class C^1 . By compactness there is $\rho_0 > 0$ so that we have for all $x \in \partial K$

$$0 < \rho_0 \leq \rho(x). \quad (7)$$

By Lemma 5 of [16] and the homothety of K and K_δ , we get that there is $R > 0$ so that we have for all $x \in \partial K$

$$K \subset B_2^n(x - R N_K(x), R). \quad (8)$$

Lemma 4 *If K is of class C^1 and homothetic to K_δ for some $\delta \leq \delta_0$, then the matrix Q is positive definite for every x_δ in ∂K_δ . Moreover δ_0 can be chosen to be*

$$\delta_0 = \frac{\rho_0^{n-1} R |B_2^{n-1}|}{n 2^{n-1}} \left(1 - \left(1 - \left(\frac{\rho_0}{4R} \right)^2 \right)^{\frac{1}{2}} \right)^n, \quad (9)$$

where ρ_0 and R are as in (7) and (8).

Proof. We want to show that we have for δ sufficiently small: For all $x \in \partial K_\delta$ and for all η we have $\cot(\beta(\eta)) > 0$. This implies that (6) is a positive-definite matrix and by Lemma 3 we get that ∂K_δ is of class C^2 and therefore ∂K is also of class C^2 by homothety.

Let ρ_0 and R be as in (7) and (8). We choose $s > 0$ so that the orthogonal projection of

$$B_2^n(x - R N_K(x), R) \cap H(x - s N_K(x), N_K(x)),$$

is contained in

$$B_2^n\left(x - t(x) N_K(x), \frac{\rho_0}{4}\right) \cap H(x - t(x) N_K(x), N_K(x)).$$




Figure 1:

It is easy to see that this holds if $s = R \left(1 - \left(1 - \left(\frac{\rho_0}{4R} \right)^2 \right)^{\frac{1}{2}} \right)$.

Now we choose δ_0 . We choose δ_0 so small that we have for all $x \in \partial K$

$$|H^-(x - s N_K(x), N_K(x)) \cap K| \geq \delta_0. \quad (10)$$

From Figure (1), it follows that

$$\cot(\alpha) \geq \frac{\frac{\rho_0}{2} - \frac{\rho_0}{4}}{t(x)} > 0.$$

From Figure (2), it follows that $\beta(\eta) \leq \gamma \leq \alpha$ and

$$\cot(\beta(\eta)) \geq \cot(\gamma) \geq \cot(\alpha) \geq \frac{\rho_0}{4 t(x)}.$$

We determine now δ_0 from (10) as follows

$$\delta_0 = \frac{\rho_0^{n-1} R |B_2^{n-1}|}{n 2^{n-1}} \left(1 - \left(1 - \left(\frac{\rho_0}{4R} \right)^2 \right)^{\frac{1}{2}} \right)^n.$$

Then for all $\delta \leq \delta_0$, (10) holds.




Figure 2:

Indeed, for $x \in \partial K$, let $C(t(x), \rho_0) = \text{conv}[x, B_2^{n-1}(x - t(x)N_K(x), \rho_0)]$ be the cone with tip at x , height $t(x)$ and with the $(n-1)$ -dimensional ball centered at $x - t(x)N_K(x)$ and radius ρ_0 as base. Then, as $B_2^n(x - t(x)N_K(x), \rho_0) \subset K$,

$$\begin{aligned}
 |H^-(x - sN_K(x), N_K(x)) \cap K| &\geq |H^-(x - sN_K(x), N_K(x)) \cap C(t(x), \rho_0)| \\
 &= \frac{\rho_0^{n-1} s^n |B_2^{n-1}|}{n t(x)^{n-1}} \geq \frac{\rho_0^{n-1} s^n |B_2^{n-1}|}{n 2^{n-1} R^{n-1}} \\
 &= \frac{\rho_0^{n-1} R |B_2^{n-1}|}{n 2^{n-1}} \left(1 - \left(1 - \left(\frac{\rho_0}{4R}\right)^2\right)^{\frac{1}{2}}\right)^n \\
 &= \delta_0 \geq \delta,
 \end{aligned}$$

which shows that (10) holds.

Remark 2 (i) Thus, by Lemmas 1, 2, 3, and 4, a convex body K in $\mathbb{R}^n$ can only be homothetic to one of its floating bodies K_δ for small enough δ , if K is strictly convex and of class C_+^2 .

(ii) The example of Remark 1 shows that the assumption of homothety is also crucial in Lemma 4.

3 Homothety conjecture for $K = B_p^n$

In this section, we show that the homothety conjecture holds true in the class of the bodies B_p^n for all $1 \leq p \leq \infty$ and their affine images.

Theorem 1 *Let $B_p^n, 1 \leq p \leq \infty$ be the unit ball of l_p^n . Let $0 < \delta < \frac{|B_p^n|}{2}$. Then $(B_p^n)_\delta = c B_p^n$ for some $0 < c < 1$ if and only if $p = 2$.*

Remark. By (1), the same holds true for affine images $T(B_p^n)$ of B_p^n under an invertible linear map T on $\mathbb{R}^n$: Let $K = T(B_p^n)$, $1 \leq p \leq \infty$. Let $0 < \delta < \frac{|K|}{2}$ be a constant. $K_\delta = c K$ for some constant $0 < c < 1$ if and only if K is an ellipsoid.

Proof of Theorem 1. It was noted above that $(B_2^n)_\delta$ is homothetic to B_2^n for all δ with $0 < \delta < \frac{|B_2^n|}{2}$.

For the reverse implication, it is enough to consider p with $1 < p < \infty$. Indeed, B_1^n and B_∞^n are polytopes and it was observed above that polytopes cannot be homothetic to any of their floating bodies.

We first consider the case when $1 < p < 2$. Then ∂B_p^n is of class C^1 but not of class C^2 : at $e_n = (0, \dots, 0, 1)$, ∂B_p^n is not C^2 . If B_p^n were homothetic to $(B_p^n)_\delta$ for some δ , then Lemma 3 would imply that $(B_p^n)_\delta$ is C^2 . Indeed, for all $x \in \partial(B_p^n)_\delta$, $0 < \beta(\eta) < \frac{\pi}{2}$ for all $\eta \in S^{n-2}$ and thus $\cot(\beta(\eta)) > 0$ or all $\eta \in S^{n-2}$. Therefore the matrix Q of (6) is positive definite and by Lemma 3 $(B_p^n)_\delta$ and thus, by homothety, B_p^n is C^2 , a contradiction.

Now we consider the case $2 \leq p < \infty$. Then ∂B_p^n is of class C^2 and thus, assuming that B_p^n is homothetic to $(B_p^n)_\delta$ for some $\delta > 0$, $(B_p^n)_\delta$ is C^2 .

It was shown in [18], that the curvature $\kappa_{B_p^n}$ at $x \in \partial B_p^n$ is

$$\kappa_{B_p^n}(x) = \frac{(p-1)^{n-1} \prod_{i=1}^n x_i^{p-2}}{(\sum_{i=1}^n |x_i|^{2(p-1)})^{\frac{1}{2}}}. \quad (11)$$

Thus for $e_n = (0, \dots, 0, 1)$, $\kappa_{B_p^n}(e_n) = 0$, if $p > 2$ and $\kappa_p(e_n) = 1$, if $p = 2$. Consequently, as we assume that $(B_p^n)_\delta = c B_p^n$ for some $\delta > 0$, the curvature at $ce_n \in \partial(B_p^n)_\delta$ is different from 0 only when $p = 2$ and for all $p > 2$

$$\kappa_{(B_p^n)_\delta}(ce_n) = 0. \quad (12)$$

By Lemma 3,

$$\kappa_{(B_p^n)_\delta}(ce_n) = \frac{1}{\det(Q)},$$

which is 0 if and only if $\det(Q) = \infty$.

By (6), the matrix Q has entries

$$Q(i, j) = \frac{1}{|K \cap H(ce_n, e_n)|} \int_{S^{n-1}} \eta_i \eta_j r^n(\eta) \cot(\beta(\eta)) d\sigma_{n-2}(\eta),$$

$1 \leq i, j \leq n-1$. Again, for all $\eta \in S^{n-2}$, $0 < \beta(\eta) < \frac{\pi}{2}$. Therefore, $\det(Q) < \infty$ and thus the curvature at $ce_n \in \partial(B_p^n)_\delta$ is strictly positive which contradicts (12).

We point out that $(B_p^n)_\delta \in C_+^2$ for all $p \in (1, \infty)$ and all $\delta > 0$ can be obtained from results in [6].

More generally, in the same way as Theorem 1, one can prove

Proposition 1 *Let K be a convex body in $\mathbb{R}^n$. If K has a point on the boundary where the Gauss curvature is either 0 or ∞ , then K is not homothetic to K_δ for any $\delta \leq \delta_0$ with δ_0 given as in Lemma 4.*

Proof. Let x_0 in ∂K be such that $\kappa_K(x_0) = \infty$ and suppose that K is homothetic to K_δ for some $\delta \leq \delta_0$. Then, by Remark 2, K is strictly convex and in C_+^2 . By Lemma 3,

$$\kappa_{K_\delta}(cx_0) = \frac{1}{\det(Q)},$$

where cx_0 is the point on ∂K_δ corresponding to x_0 by homothety. This is ∞ if and only if $\det(Q) = 0$. But by Lemmas 3 and 4 (respectively Remark 2), K is in C_+^2 , thus Q is positive definite and thus $\det(Q) > 0$, a contradiction.

The case $\kappa_K(x_0) = 0$ is treated similarly.

4 Homothety conjecture for general K

In Section 2 we proved the homothety conjecture for B_p^n , $1 \leq p \leq \infty$ and their affine images. The proof uses the fact that one only needs to examine one properly chosen direction in order to be able to conclude. In this section, we will use two directions to prove positive solutions to the homothety conjecture. Moreover, our approach

is robust and can be used to obtain (partial) positive solutions for *generalized homothety conjecture*.

Theorem 2 *Let K be a convex body in $\mathbb{R}^n$. There exists a positive number $\delta(K)$, such that K_δ is homothetic to K for some $\delta < \delta(K)$, if and only if K is an ellipsoid.*

In the next section we provide estimates for the threshold $\delta(K)$.

To prove this theorem, we need the following lemmas. Lemma 5 was proved in [10] and Lemma 6 was proved in [15]. See also [26] for similar results.

Lemma 5 [10] *Let $K \in C_+^2$ be a convex body with boundary of class C^2 and everywhere strictly positive Gaussian curvature. Let $c(K, n)$ be a constant only depending on K and n . Then*

$$\kappa_K(x) \langle x, N_K(x) \rangle^{-(n+1)} = c(K, n), \quad \forall x \in \partial K$$

holds true if and only if K is an ellipsoid.

Lemma 6 [15] *Let $K \in C_+^2$ be a convex body with boundary of class C^2 and everywhere strictly positive Gaussian curvature. Then, for any $x \in \partial K$,*

$$\lim_{\delta \rightarrow 0} c_n \frac{\langle x, N_K(x) \rangle}{n \delta^{\frac{2}{n+1}}} \left[1 - \left(\frac{\|x_\delta\|}{\|x\|} \right)^n \right] = (\kappa_K(x))^{\frac{1}{n+1}},$$

where $x_\delta \in \partial(K_\delta) \cap [0, x]$ and $c_n = 2 \left(\frac{|B_2^{n-1}|}{n+1} \right)^{\frac{2}{n+1}}$.

Proof of Theorem 2. Suppose that K is homothetic to K_δ for some $\delta < \delta(K)$ with $\delta(K) \leq \delta_0$, where δ_0 is given by Lemma 4 ($\delta(K)$ will be determined more precisely later). Suppose that K not an ellipsoid. By Lemma 1 and homothety, K is strictly convex. By Lemma 2 and homothety, K is C^1 . By Lemma 4, homothety and Lemma 3, K is in C_+^2 . $K \in C_+^2$ implies that $\langle x, N_K(x) \rangle$ and $\kappa_K(x)$ both are continuous functions on ∂K . We define points $x_M \in \partial K$ and $x_m \in \partial K$ by

$$\begin{aligned} (T_M)^{n+1} &= \frac{\kappa_K(x_M)}{\langle x_M, N_K(x_M) \rangle^{n+1}} = \max_{x \in \partial K} \frac{\kappa_K(x)}{\langle x, N_K(x) \rangle^{n+1}}, \\ (T_m)^{n+1} &= \frac{\kappa_K(x_m)}{\langle x_m, N_K(x_m) \rangle^{n+1}} = \min_{x \in \partial K} \frac{\kappa_K(x)}{\langle x, N_K(x) \rangle^{n+1}}. \end{aligned}$$

As K is not an ellipsoid, Lemma 5 implies that

$$\frac{\kappa_K(x_M)}{\langle x_M, N_K(x_M) \rangle^{n+1}} > \frac{\kappa_K(x_m)}{\langle x_m, N_K(x_m) \rangle^{n+1}}.$$

Equivalently, one has

$$\tau = \frac{T_M}{T_m} > 1.$$

By Lemma 6, for $\varepsilon_1 = \frac{\tau-1}{3\tau}$, there exists $\delta_1(K) > 0$, such that, for all $\delta \leq \delta_1(K)$,

$$\begin{aligned} \frac{c_n}{n \delta^{\frac{2}{n+1}}} \left[1 - \left(\frac{\|x_{M,\delta}\|}{\|x_M\|} \right)^n \right] &\geq \frac{(\kappa_K(x_M))^{\frac{1}{n+1}}}{\langle x_M, N_K(x_M) \rangle} (1 - \varepsilon_1) \\ &= T_M (1 - \varepsilon_1) = T_M \left(\frac{2\tau + 1}{3\tau} \right), \end{aligned} \quad (13)$$

where $x_{M,\delta} = \partial K_\delta \cap [0, x_M]$.

Again by Lemma 6, for $\varepsilon_2 = \frac{\tau-1}{2}$, there exists $\delta_2(K) > 0$, such that, for all $\delta \leq \delta_2(K)$,

$$\begin{aligned} \frac{c_n}{n \delta^{\frac{2}{n+1}}} \left[1 - \left(\frac{\|x_{m,\delta}\|}{\|x_m\|} \right)^n \right] &\leq \frac{(\kappa_K(x_m))^{\frac{1}{n+1}}}{\langle x_m, N_K(x_m) \rangle} (1 + \varepsilon_2) \\ &= T_m (1 + \varepsilon_2) = T_m \left(\frac{\tau + 1}{2} \right), \end{aligned} \quad (14)$$

where $x_{m,\delta} = \partial K_\delta \cap [0, x_m]$.

Let $\delta(K) = \min\{\delta_0, \delta_1(K), \delta_2(K)\}$. If $K_\delta = cK$ for some $0 < \delta < \delta(K)$ and for some constant $c(\delta) > 0$, then formulas (13) and (14) imply that

$$T_M \left(\frac{2\tau + 1}{3\tau} \right) \leq \frac{c_n (1 - c^n)}{n \delta^{\frac{2}{n+1}}} \leq T_m \left(\frac{\tau + 1}{2} \right),$$

and hence

$$T_M \left(\frac{2\tau + 1}{3\tau} \right) \leq T_m \left(\frac{\tau + 1}{2} \right).$$

Equivalently, $\tau \leq 1$ which is a contradiction. Hence K must be an ellipsoid.

Remark. We can replace x_M and x_m by any two points $x, y \in \partial K$ such that

$$\frac{\kappa_K(x)}{\langle x, N_K(x) \rangle^{n+1}} \neq \frac{\kappa_K(y)}{\langle y, N_K(y) \rangle^{n+1}}.$$

Following the above proof, one gets

Theorem 3 *Let K be a convex body in $\mathbb{R}^n$. Suppose that there are two points $x, y \in \partial K$, such that, both $\kappa_K(x)$ and $\kappa_K(y)$ exist and are finite, and*

$$\kappa_K(x)\langle x, N_K(x) \rangle^{-(n+1)} \neq \kappa_K(y)\langle y, N_K(y) \rangle^{-(n+1)}.$$

Then there is a constant $\delta(K, x, y)$ depending on K, x and y such that K_δ is not homothetic to K for all $\delta \leq \delta(K, x, y)$.

Analogously, we can ask the following *generalized homothety conjecture*.

Generalized Homothety Conjecture: *Let K be a convex body in $\mathbb{R}^n$. Does $K_s = cK$ for some $0 < s$ and some $c > 0$ imply that K is an ellipsoid? Here $\{K_s\}_{s \geq 0}$ is a family of convex bodies constructed from K .*

Besides the convex floating body K_δ , examples of such K_s include

1. The illumination body K^δ [22]

$$K^\delta = \{x \in \mathbb{R}^n : |\text{conv}(x, K)| - |K| \leq \delta\}.$$

2. The convolution body $C(K, t)$ [3, 12] which, for a symmetric convex body K in $\mathbb{R}^n$ and $t \geq 0$ is defined by

$$C(K, t) = \left\{ \frac{x}{2} \in \mathbb{R}^n : |K \cap (K + x)| \geq 2t \right\}.$$

3. The Santaló-regions $S(K, t)$ [7]

$$S(K, t) = \left\{ x \in K : |K^x| \leq \frac{1}{t} \right\},$$

where $K^x = (K - x)^\circ = \{z \in \mathbb{R}^n : \langle z, y - x \rangle \leq 1, \forall y \in K\}$ is the polar body of K with respect to $x \in K$.

We refer to [23, 24, 25] for more general constructions.

The following theorem provides (partial) positive solutions of the generalized homothety conjecture. Theorem 4 (i) was proved with a different method in [20].

Theorem 4 *Let K be a convex body in C_+^2 .*

(i) [20] There exists a positive number $\tilde{\delta}(K)$ such that K^δ is homothetic to K for some $\delta < \tilde{\delta}(K)$, if and only if K is an ellipsoid.

(ii) There exists a positive number $t(K)$ such that $C(K, t)$ is homothetic to K for some $t < t(K)$, if and only if K is an ellipsoid.

(iii) There exists a positive number $\tilde{t}(K)$ such that $S(K, t)$ is homothetic to K for some $t < \tilde{t}(K)$, if and only if K is an ellipsoid.

Remark. The proof of this theorem is same as the proof of Theorem 2. The proof of (i) also relies on Lemma 3 in [22]. For the proof of (ii), we refer to results similar to Lemma 6 in [12]. For (iii), one uses Lemma 13 in [7].

Estimates on the thresholds $\tilde{\delta}(K)$, $t(K)$ and $\tilde{t}(K)$ can be obtained similar to the one for $\delta(K)$. This is treated in the next section.

5 Estimates on the threshold $\delta(K)$

Our proof of Theorem 2 gives a possibility to estimate the threshold $\delta(K)$ for a convex body K in $\mathbb{R}^n$. Thus we assume that K_δ is homothetic to K for some $\delta \leq \delta(K)$ with 0 as the center of homothety. Let $x_M \in \partial K$, $x_m \in \partial K$, T_M , T_m and τ be as in the previous section. Note that the points x_m, x_M may not be uniquely determined. We just choose any two points satisfying the condition. Let

$$r_m = \kappa_K(x_m)^{-\frac{1}{n-1}} \quad \text{and} \quad r_M = \kappa_K(x_M)^{-\frac{1}{n-1}}.$$

Let

$$a = \min \left\{ 1 - \left(\frac{2}{1+\tau} \right)^{\frac{n+1}{n-1}}, \left(\frac{3\tau}{1+2\tau} \right)^{\frac{n+1}{n-1}} - 1 \right\}. \quad (15)$$

We show

Theorem 5 *Let K be a convex body in $\mathbb{R}^n$ with everywhere on ∂K strictly positive Gauss curvature and such that ∂K is C^3 . Let a be as in (15). Then $\delta(K)$ of Theorem 2 can be chosen to be*

$$\delta(K) = \min \left\{ \delta_0, \delta_1, \delta_2, \delta_m, \delta_M, \frac{(1-a)^n r_m^n |B_2^n|}{2}, \frac{(1-a)^n r_M^n |B_2^n|}{2} \right\},$$

where δ_0 is as in (9), and the expressions of $\delta_1, \delta_2, \delta_m, \delta_M$ are in the proof.

Proof. For $\alpha = (\alpha_1, \dots, \alpha_m) \in \mathbb{N}^m$, and an $|\alpha|$ -times continuously differentiable function g , let

$$\begin{aligned} |\alpha| &= \alpha_1 + \dots + \alpha_m, \\ \alpha! &= \alpha_1! \dots \alpha_m!, \\ D^\alpha g &= D_1^{\alpha_1} \dots D_n^{\alpha_m} g = \frac{\partial^{|\alpha|} g}{\partial t_1^{\alpha_1} \dots \partial t_n^{\alpha_m}}, \end{aligned}$$

where $D_i^{\alpha_i} = D_j \dots D_j$ is the α_i times product of D_i .

As determining $\delta(K)$ is invariant under affine transformations of determinant 1, we can assume that the ellipsoid approximating ∂K at x_m is a Euclidean ball and then have (see [17]): For $a > 0$ given as in (15) above, there exists $\Delta_{a,m}$ such that for all $\Delta \leq \Delta_{a,m}$

$$\begin{aligned} B_2^n(x_m - \bar{r}_m N_K(x_m), \bar{r}_m) \cap H^-(x_m - \Delta N_K(x_m), N_K(x_m)) \\ \subseteq K \cap H^-(x_m - \Delta N_K(x_m), N_K(x_m)), \end{aligned} \quad (16)$$

where $\bar{r}_m = (1 - a)r_m$. In addition, we also choose $\Delta_{a,m} < \bar{r}_m$.

Assume now that $x_m = 0$, that $N_K(x_m) = -e_n$ and that the other $(n - 1)$ axes of the approximating ellipsoid coincide with the remaining $(n - 1)$ coordinate axes. Locally we can then describe ∂K by a convex function $f_m : \mathbb{R}^{n-1} \rightarrow \mathbb{R}$, such that $(t_1, \dots, t_{n-1}) \rightarrow (t_1, \dots, t_{n-1}, f_m(t_1, \dots, t_{n-1})) \in \partial K$.

As ∂K is C^3 , by Taylor's theorem there exists $s \in P(K)$ such that for all $t = (t_1, \dots, t_{n-1}) \in P(K)$, the orthogonal projection of K onto $\mathbb{R}^{n-1}$,

$$f_m(t) = \frac{1}{2} \langle t, At \rangle + \sum_{|\alpha|=3} \frac{D^\alpha f_m(s)}{\alpha!} t^\alpha,$$

where $t^\alpha = t_1^{\alpha_1} \dots t_{n-1}^{\alpha_{n-1}}$ and $A = \left(\frac{\partial^2 f_m}{\partial t_i \partial t_j}(0) \right)_{i,j=1}^{n-1}$ is the Hessian of f_m at 0. Clearly, $A = \frac{1}{r_m} Id_{n-1}$ with Id_{n-1} the identity matrix. $K \in C^3$ also implies that $|D^\alpha f_m(t)| \leq D$ for some $D > 0$, for all α with $|\alpha| = 3$, and all $t \in P(K)$. Therefore,

$$f_m(t) \leq \frac{1}{2} \langle t, At \rangle + \frac{(n-1)^3}{6} D \|t\|^3 = \frac{1}{2} \frac{\|t\|^2}{r_m} + \frac{(n-1)^3}{6} D \|t\|^3. \quad (17)$$

$\partial B_2^n(x_m - \bar{r}_m, \bar{r}_m)$ is described in our chosen coordinate system by a function

$g_m : \mathbb{R}^{n-1} \rightarrow \mathbb{R}$ - for $t_n = g_m(t) \leq \bar{r}_m$ -

$$t_n = g_m(t) = \bar{r}_m \left[1 - \sqrt{1 - \left(\frac{\|t\|}{\bar{r}_m} \right)^2} \right], \quad \text{if } t_n \leq \bar{r}_m.$$

As $\frac{b}{2} \leq 1 - \sqrt{1-b}$ for all $b \in [0, 1]$, we get for all t with $\|t\| \leq \bar{r}_m$

$$t_n = g_m(t) \geq \frac{1}{2} \frac{\|t\|^2}{(1-a)r_m}. \quad (18)$$

Thus by (17) and (18), for

$$\|t\| \leq t_{a,m} = \min \left\{ \bar{r}_m, \frac{3a}{D\bar{r}_m(n-1)^3} \right\},$$

one has $t_n = g_m(t) \geq f_m(t)$. We then let

$$\Delta_{a,m} = \bar{r}_m - \sqrt{\bar{r}_m^2 - t_{a,m}^2},$$

and hence for all $\Delta \leq \Delta_{a,m}$, condition (16) holds true. We further let

$$\delta_{m,1} = |B_2^{n-1}| \int_{\bar{r}_m - \Delta_{a,m}}^{\bar{r}_m} (\bar{r}_m^2 - y^2)^{\frac{n-1}{2}} dy. \quad (19)$$

Denote by $\Delta_{\bar{r}_m}$ the height of a cap of $\bar{r}_m B_2^n$ of volume exactly δ . Recall $x_{m,\delta} = \partial K_\delta \cap [0, x_m]$, and

$$\Delta_{x_m} = \left\langle \frac{x_m}{\|x_m\|}, N_K(x_m) \right\rangle \|x_m - x_{m,\delta}\|.$$

By (16) and (19), one has, for all $\delta \leq \delta_{m,1}$,

$$\Delta_{x_m} \leq \Delta_{\bar{r}_m}.$$

Moreover, as x and $x_{m,\delta}$ are colinear,

$$1 - \left(\frac{\|x_{m,\delta}\|}{\|x_m\|} \right)^n \leq n \frac{\|x_m - x_{m,\delta}\|}{\|x_m\|}.$$

Thus

$$\frac{c_n}{\delta^{\frac{2}{n+1}}} \frac{\|x_m - x_{m,\delta}\|}{\|x_m\|} = \frac{c_n \Delta_{x_m}}{\delta^{\frac{2}{n+1}} \langle x_m, N_K(x_m) \rangle} \leq \frac{c_n \Delta_{\bar{r}_m}}{\delta^{\frac{2}{n+1}} \langle x_m, N_K(x_m) \rangle}. \quad (20)$$

Let $\delta < \frac{|B_2^n| \bar{r}_m^n}{2}$. Then $\Delta_{\bar{r}_m} < \bar{r}_m$ and by definition of $\Delta_{\bar{r}_m}$,

$$\begin{aligned} \delta &= |B_2^{n-1}| \int_{\bar{r}_m - \Delta_{\bar{r}_m}}^{\bar{r}_m} (\bar{r}_m - y)^{\frac{n-1}{2}} (\bar{r}_m + y)^{\frac{n-1}{2}} dy \\ &\geq |B_2^{n-1}| (2\bar{r}_m - \Delta_{\bar{r}_m})^{\frac{n-1}{2}} \int_{\bar{r}_m - \Delta_{\bar{r}_m}}^{\bar{r}_m} (\bar{r}_m - y)^{\frac{n-1}{2}} dy \\ &= 2^{\frac{n+1}{2}} \bar{r}_m^{\frac{n-1}{2}} \frac{|B_2^{n-1}|}{n+1} \left(1 - \frac{\Delta_{\bar{r}_m}}{2\bar{r}_m}\right)^{\frac{n-1}{2}} \Delta_{\bar{r}_m}^{\frac{n+1}{2}}. \end{aligned}$$

In particular, using $\left(1 - \frac{\Delta_{a,m}}{2\bar{r}_m}\right) > \frac{1}{2}$ and $\Delta_{a,m} = \bar{r}_m \left(1 - \sqrt{1 - \frac{t_{a,m}^2}{\bar{r}_m^2}}\right) \geq \frac{t_{a,m}^2}{2\bar{r}_m}$,

$$\delta_{m,1} \geq \frac{t_{a,m}^{n+1} |B_2^{n-1}|}{2^{\frac{n-1}{2}} (n+1) \bar{r}_m} = \delta_m. \quad (21)$$

And

$$\begin{aligned} \delta &= |B_2^{n-1}| \int_{\bar{r}_m - \Delta_{\bar{r}_m}}^{\bar{r}_m} (\bar{r}_m - y)^{\frac{n-1}{2}} (\bar{r}_m + y)^{\frac{n-1}{2}} dy \\ &\leq |B_2^{n-1}| 2^{\frac{n-1}{2}} \bar{r}_m^{\frac{n-1}{2}} \int_{\bar{r}_m - \Delta_{\bar{r}_m}}^{\bar{r}_m} (\bar{r}_m - y)^{\frac{n-1}{2}} dy = 2^{\frac{n+1}{2}} \bar{r}_m^{\frac{n-1}{2}} \frac{|B_2^{n-1}|}{n+1} \Delta_{\bar{r}_m}^{\frac{n+1}{2}}. \end{aligned}$$

Hence for all $\delta < \frac{\bar{r}_m^n |B_2^n|}{2}$,

$$\frac{\bar{r}_m^{-\frac{n-1}{n+1}}}{c_n} \leq \frac{\Delta_{\bar{r}_m}}{\delta^{\frac{2}{n+1}}} \leq \frac{\bar{r}_m^{-\frac{n-1}{n+1}}}{c_n} \left(1 - \frac{\Delta_{\bar{r}_m}}{2\bar{r}_m}\right)^{-\frac{n-1}{n+1}}. \quad (22)$$

From (20), (21), and (22) we get for all $\delta < \min\{\frac{|B_2^n| \bar{r}_m^n}{2}, \delta_m\}$,

$$\begin{aligned} \frac{c_n}{n \delta^{\frac{2}{n+1}}} \left[1 - \left(\frac{\|x_{m,\delta}\|}{\|x_m\|}\right)^n\right] &\leq \frac{\bar{r}_m^{-\frac{n-1}{n+1}}}{\langle x_m, N_K(x_m) \rangle} \left(1 - \frac{\Delta_{\bar{r}_m}}{2\bar{r}_m}\right)^{-\frac{n-1}{n+1}} \\ &= (1-a)^{-\frac{n-1}{n+1}} \frac{\kappa_K(x_m)^{\frac{1}{n+1}}}{\langle x_m, N_K(x_m) \rangle} \left(1 - \frac{\Delta_{\bar{r}_m}}{2\bar{r}_m}\right)^{-\frac{n-1}{n+1}} \\ &= T_m (1-a)^{-\frac{n-1}{n+1}} \left(1 - \frac{\Delta_{\bar{r}_m}}{2\bar{r}_m}\right)^{-\frac{n-1}{n+1}}. \end{aligned}$$

Hence for all $\delta \leq \min\{\delta_2, \frac{|B_2^n| \bar{r}_m^n}{2}, \delta_m\}$, (14) of the previous section will hold, if we choose δ_2 so that

$$\left(1 - \frac{\Delta_{\bar{r}_m}}{2\bar{r}_m}\right)^{-\frac{n-1}{n+1}} \leq (1-a)^{\frac{n-1}{n+1}} \left(\frac{\tau+1}{2}\right). \quad (23)$$

By (22) and (23),

$$\Delta_{\bar{r}_m} \leq \delta^{\frac{2}{n+1}} \frac{\bar{r}_m^{-\frac{n-1}{n+1}}}{c_n} \left(1 - \frac{\Delta_{\bar{r}_m}}{2\bar{r}_m}\right)^{-\frac{n-1}{n+1}} \leq \delta^{\frac{2}{n+1}} \frac{\bar{r}_m^{-\frac{n-1}{n+1}}}{c_n} (1-a)^{\frac{n-1}{n+1}} \left(\frac{\tau+1}{2}\right).$$

Thus, to have (23), it is enough to have that

$$\left(1 - \frac{\delta^{\frac{2}{n+1}}}{c_n \bar{r}_m^{\frac{2n}{n+1}}} \left(\frac{\tau+1}{4}\right) (1-a)^{\frac{n-1}{n+1}}\right)^{-1} \leq (1-a) \left(\frac{\tau+1}{2}\right)^{\frac{n+1}{n-1}}.$$

Hence, we can let

$$\delta_2 = 2^{\frac{3(n+1)}{2}} \left(\frac{1-a}{1+\tau}\right)^{\frac{n+1}{2}} r_m^n \frac{|B_2^{n-1}|}{n+1} \left[1 - (1-a)^{-1} \left(\frac{2}{\tau+1}\right)^{\frac{n+1}{n-1}}\right]^{\frac{n+1}{2}}. \quad (24)$$

Now we consider x_M . We let $r_M = \kappa_K(x_M)^{\frac{1}{1-n}}$, $\bar{r}_M = (1-a)r_M$ and $\bar{R}_M = (1+a)r_M$. Assume now that $x_M = 0$, that $N_K(x_M) = -e_n$ and that the other $(n-1)$ axes of the approximating ellipsoid coincide with the remaining $(n-1)$ coordinate axes. Locally we can then describe ∂K by a convex function $f_M : \mathbb{R}^{n-1} \rightarrow \mathbb{R}$, such that $(t_1, \dots, t_{n-1}) \rightarrow (t_1, \dots, t_{n-1}, f_M(t_1, \dots, t_{n-1})) \in \partial K$.

For

$$t_{M,1} = \min \left\{ \bar{r}_M, \frac{3a}{D\bar{r}_M(n-1)^3} \right\},$$

let

$$\Delta_{M,1} = \bar{r}_M - \sqrt{\bar{r}_M^2 - t_{M,1}^2}.$$

We repeat the previous argument and get for all $\Delta \leq \Delta_{M,1}$, that

$$\begin{aligned} B_2^n(x_M - \bar{r}_M N_K(x_M), \bar{r}_M) \cap H^-(x_M - \Delta N_K(x_M), N_K(x_M)) \\ \subseteq K \cap H^-(x_M - \Delta N_K(x_M), N_K(x_M)). \end{aligned} \quad (25)$$

Let $\xi = 1 + \frac{a}{2}$. Then for all $b \in [0, \frac{4(\xi-1)}{\xi^2}]$,

$$1 - \sqrt{1-b} \leq \frac{\xi}{2} b.$$

For $g_M(t) \leq \bar{R}_M$, let

$$t_n = g_M(t) = \bar{R}_M \left[1 - \sqrt{1 - \left(\frac{\|t\|}{\bar{R}_M}\right)^2}\right],$$

be the function that describes $\partial B_2^n(x_M - \bar{R}_M N_K(x_M), \bar{R}_M)$. For $\|(t_1, \dots, t_{n-1})\| \leq \frac{2(\xi-1)^{\frac{1}{2}}}{\xi} \bar{R}_M$, we get that

$$t_n = g_M(t) \leq \frac{\xi}{2} \frac{\|t\|^2}{\bar{R}_M}.$$

As in inequality (17),

$$f_M(t) \geq \frac{1}{2} \frac{\|t\|^2}{r_M} - \frac{(n-1)^3}{6} D \|t\|^3.$$

Therefore, for $\|t\| \leq t_{M,2}$ where

$$t_{M,2} = \min \left\{ \frac{2(\xi-1)^{\frac{1}{2}}}{\xi} \bar{R}_M, \frac{3a}{2D\bar{R}_M(n-1)^3} \right\},$$

we get that $f_M(t) \geq g_M(t) = t_n$ which implies that

$$\begin{aligned} B_2^n(x_M - \bar{R}_M N_K(x_M), \bar{R}_M) \cap H^-(x_M - \Delta N_K(x_M), N_K(x_M)) \\ \supseteq K \cap H^-(x_M - \Delta N_K(x_M), N_K(x_M)), \end{aligned} \quad (26)$$

for all $\Delta \leq \Delta_{M,2}$ with $\Delta_{M,2} = \bar{R}_M - \sqrt{\bar{R}_M^2 - t_{M,2}^2}$.

We let $\Delta_{a,M} = \min\{\Delta_{M,1}, \Delta_{M,2}\}$, and

$$\delta_{M,1} = |B_2^{n-1}| \int_{\bar{r}_M - \Delta_{a,M}}^{\bar{r}_M} (\bar{r}_M^2 - y^2)^{\frac{n-1}{2}} dy. \quad (27)$$

In particular, as above,

$$\delta_{M,1} \geq 2 \frac{|B_2^{n-1}|}{n+1} \bar{r}_M^{\frac{n-1}{2}} \Delta_{a,M}^{\frac{n+1}{2}} = \delta_M, \quad (28)$$

where $\Delta_{a,M}$ can be taken as

$$\min \left\{ \frac{t_{M,1}^2}{2(1-a)r_m}, \frac{t_{M,2}^2}{2(1-a)\bar{R}_M} \right\}.$$

Let $\Delta_{\bar{R}_M}$ be the height of a cap of $\bar{R}_M B_2^n$ of volume exactly δ . By (25), (26), (27), and (28), for all $\delta \leq \delta_M$,

$$\Delta_{\bar{R}_M} \leq \Delta_{x_M} = \left\langle \frac{x_M}{\|x_M\|}, N_K(x_M) \right\rangle \|x_M - x_{M,\delta}\|.$$

Recall $x_{M,\delta} = [0, x_M] \cap \partial K_\delta$. Similar to (22), for all $\delta < \min\{\frac{\bar{r}_M^n |B_2^n|}{2}, \delta_M\}$,

$$\frac{\bar{R}_M^{-\frac{n-1}{n+1}}}{c_n} \leq \frac{\Delta_{\bar{R}_M}}{\delta^{\frac{2}{n+1}}} \leq \frac{\bar{R}_M^{-\frac{n-1}{n+1}}}{c_n} \left(1 - \frac{\Delta_{\bar{R}_M}}{2\bar{R}_M}\right)^{-\frac{n-1}{n+1}}.$$

This implies that, for all $\delta < \min\{\frac{\bar{r}_M^n |B_2^n|}{2}, \delta_M\}$,

$$\begin{aligned}
\frac{c_n}{n \delta^{\frac{2}{n+1}}} \left[1 - \left(\frac{\|x_{M,\delta}\|}{\|x_M\|} \right)^n \right] &= \frac{c_n}{n \delta^{\frac{2}{n+1}}} \left[1 - \left(1 - \frac{\Delta_{x_M}}{\langle x_M, N_K(x_M) \rangle} \right)^n \right] \\
&\geq \frac{c_n}{n \delta^{\frac{2}{n+1}}} \left[1 - \left(1 - \frac{\Delta_{\bar{R}_M}}{\langle x_M, N_K(x_M) \rangle} \right)^n \right] \\
&\geq \frac{c_n}{\delta^{\frac{2}{n+1}}} \left(\frac{\Delta_{\bar{R}_M}}{\langle x_M, N_K(x_M) \rangle} - \frac{(n-1) \Delta_{\bar{R}_M}^2}{2 \langle x_M, N_K(x_M) \rangle^2} \right) \\
&\geq \frac{\bar{R}_M^{-\frac{n-1}{n+1}}}{\langle x_M, N_K(x_M) \rangle} \left(1 - \frac{(n-1) \Delta_{\bar{R}_M}}{2 \langle x_M, N_K(x_M) \rangle} \right) \\
&\geq (1+a)^{-\frac{n-1}{n+1}} T_M \left(1 - \frac{(n-1) \delta^{\frac{2}{n+1}} T_M}{2 c_n (1+a)^{\frac{n-1}{n+1}}} \left[1 - \frac{\Delta_{\bar{R}_M}}{2 \bar{R}_M} \right]^{-\frac{n-1}{n+1}} \right) \\
&\geq (1+a)^{-\frac{n-1}{n+1}} T_M \left(1 - \frac{(n-1) T_M \delta^{\frac{2}{n+1}}}{2^{\frac{2}{n+1}} c_n (1+a)^{\frac{n-1}{n+1}}} \right).
\end{aligned}$$

The last inequality holds as $\frac{\Delta_{\bar{R}_M}}{2 \bar{R}_M} \leq \frac{1}{2}$. To have (13) of the previous section, it is enough to have

$$(1+a)^{-\frac{n-1}{n+1}} \left(1 - \frac{(n-1) T_M \delta^{\frac{2}{n+1}}}{2^{\frac{2}{n+1}} c_n (1+a)^{\frac{n-1}{n+1}}} \right) \geq \frac{2\tau+1}{3\tau},$$

or, equivalently,

$$\delta \leq \frac{[1 - (1+a)^{\frac{n-1}{n+1}} (\frac{2\tau+1}{3\tau})]^{\frac{n+1}{2}} 2^{\frac{n+3}{2}} |B_2^{n-1}| (1+a)^{\frac{n-1}{2}}}{(n-1)^{\frac{n+1}{2}} T_M^{\frac{n+1}{2}} (n+1)} (=:\delta_1). \quad (29)$$

Now we let the threshold $\delta(K)$ be

$$\delta(K) = \min \left\{ \delta_0, \delta_1, \delta_2, \delta_m, \delta_M, \frac{(1-a)^n r_m^n |B_2^n|}{2}, \frac{(1-a)^n r_M^n |B_2^n|}{2} \right\},$$

where $\delta_0, \delta_1, \delta_2, \delta_m, \delta_M$ are as in (9), (29), (24), (21) and (28) respectively.

References

- [1] I. Barany and R. A. Vitale, *Random convex hulls: Floating bodies and expectations*. J. Approx. Theory, **75** (1993) 130-135.

- [2] C. Buchta and M. Reitzner, *Equiaffine inner parallel curves of a plane convex body and the convex hulls of randomly chosen points*. Probab. Theory and Related Fields, **108** (1997) 385-415.
- [3] K. Kiener, *Extremalität von Ellipsoiden und die Faltungsungleichung von Sobolev*. Arch. Math. **46** (1986) 162-168.
- [4] K. Leichtweiss, *Über Eine Formel Blaschkes Zur Affinoberfläche*. Studia Scientiarum Mathematicarum Hungarica, **21** (1986) 453-474.
- [5] M. Meyer and S. Reisner, *Characterization of ellipsoids by section-centroid location*. Geometriae Dedicata, **31** (1989) 345-355.
- [6] M. Meyer and S. Reisner, *A Geometric Property of the Boundary of Symmetric Convex Bodies and Convexity of Flotation Surface*. Geometriae Dedicata, **37** (1991) 327-337.
- [7] M. Meyer and E. Werner, *The Santaló-regions of a convex body*. Trans. Amer. Math. Soc. **350** (11) (1998) 4569-4591.
- [8] M. Meyer and E. Werner, *On the p -affine surface area*. Adv. Math. **152** (2000) 288-313.
- [9] V. D. Milman and A. Pajor, *Isotropic position and inertia ellipsoids and zonoids of the unit ball of an n -dimensional normed space*. Israel Seminar on GAFA, Lindenstrauss, Milman (Eds.). Springer Lect. Notes in Math. **1376** (1989) 107-131.
- [10] C. M. Petty, *Affine isoperimetric problems*. Ann. New York Acad. Sci. **440** (1985) 113-127.
- [11] R. T. Rockafellar, *Convex Analysis*. Princeton University Press, 1970.
- [12] M. SCHMUCKENSCHLÄGER, *The distribution function of the convolution square of a convex symmetric body in $\mathbb{R}^n$* . Israel Journal of Math. **78** (1992) 309-334.
- [13] C. Schütt, *The convex floating body and polyhedral approximation*. Israel J. Math. **73** (1991) 65-77.
- [14] C. Schütt, *Floating Body, Illumination Body, and Polytopal Approximation*. Convex Geometric Analysis, MSRI Publications, **34** (1998) 203-229.
- [15] C. Schütt and E. Werner, *The convex floating body*. Math. Scand. **66** (1990) 275-290.
- [16] C. Schütt and E. Werner, *Homothetic floating bodies*. Geom. Dedicata, **49** (1994) 335-348.

- [17] C. Schütt and E. Werner, *Random polytopes of points chosen from the boundary of a convex body*, in: GAFA Seminar Notes, in: Lecture Notes in Math., vol. 1807, Springer-Verlag, 2002, pp. 241-422.
- [18] C. Schütt and E. Werner, *Surface bodies and p -affine surface area*. Adv. Math. **187** (2004) 98-145.
- [19] A. Stancu, *The Floating Body Problem*. Bull. London Math. Soc. **38** (2006) 839-846.
- [20] A. Stancu, *Two volume product inequalities and their applications*. To appear in Canadian Math. Bull.
- [21] A. Stancu and E. Werner, *New Higher Order Equiaffine Invariants*. Israel J. Math. **171** (2009), 221-235.
- [22] E. Werner, *Illumination bodies and affine surface area*. Studia Math. **110** (1994) 257-269.
- [23] E. Werner, *A general geometric construction for affine surface area*. Studia Math. **132** (1999) 227-238.
- [24] E. Werner, *The p -affine surface area and geometric interpretations*. Rend. Circ. Math. Palermo, **70** (2002) 367-382.
- [25] E. Werner, *On L_p affine surface areas*. Indiana Univ. Math. J. **56** (2007) 2305-2324
- [26] E. Werner and D. Ye, *New L_p -affine isoperimetric inequalities*. Adv. Math. **218** (2008) 762-780.

Elisabeth Werner

Department of Mathematics

Case Western Reserve University

Cleveland, Ohio 44106, U. S. A.

`elisabeth.werner@case.edu`

Université de Lille 1

UFR de Mathématique

59655 Villeneuve d'Ascq, France

Deping Ye

Department of Mathematics

202 Mathematical Sciences Bldg

University of Missouri

Columbia, MO 65211 USA

`yed@missouri.edu`